
BCL6 enables Ph(+) acute lymphoblastic leukaemia cells to survive BCR-ABL1 kinase inhibition.

Journal: Nature

Publication Year: 2011

Authors: Cihangir Duy, Christian Hurtz, Seyedmehdi Shojaee, Leandro Cerchietti, Huimin Geng, Srividya Swaminathan, Lars Klemm, Soo-Mi Kweon, Rahul Nahar, Melanie Braig, Eugene Park, Yong-Mi Kim, Wolf-Karsten Hofmann, Sebastian Herzog, Hassan Jumaa, H Phillip Koeffler, J Jessica Yu, Nora Heisterkamp, Thomas G Graeber, Hong Wu, B Hilda Ye, Ari Melnick, Markus Muschen

PubMed link: 21593872

Funding Grants: Dual targeting of tyrosine kinase and BCL6 signaling for leukemia stem cell eradication

Public Summary:

Tyrosine kinase inhibitors (TKIs) are widely used to treat patients with leukaemia driven by BCR-ABL1 (ref. 1) and other oncogenic tyrosine kinases. Recent efforts have focused on developing more potent TKIs that also inhibit mutant tyrosine kinases. However, even effective TKIs typically fail to eradicate leukaemia-initiating cells (LICs), which often cause recurrence of leukaemia after initially successful treatment. Here we report the discovery of a novel mechanism of drug resistance, which is based on protective feedback signalling of leukaemia cells in response to treatment with TKI. We identify BCL6 as a central component of this drug-resistance pathway and demonstrate that targeted inhibition of BCL6 leads to eradication of drug-resistant and leukaemia-initiating subclones.

Scientific Abstract:

Tyrosine kinase inhibitors (TKIs) are widely used to treat patients with leukaemia driven by BCR-ABL1 (ref. 1) and other oncogenic tyrosine kinases. Recent efforts have focused on developing more potent TKIs that also inhibit mutant tyrosine kinases. However, even effective TKIs typically fail to eradicate leukaemia-initiating cells (LICs), which often cause recurrence of leukaemia after initially successful treatment. Here we report the discovery of a novel mechanism of drug resistance, which is based on protective feedback signalling of leukaemia cells in response to treatment with TKI. We identify BCL6 as a central component of this drug-resistance pathway and demonstrate that targeted inhibition of BCL6 leads to eradication of drug-resistant and leukaemia-initiating subclones.

Source URL: <https://www.cirm.ca.gov/about-cirm/publications/bcl6-enables-ph-acute-lymphoblastic-leukaemia-cells-survive-bcr-abl1-kinase>